

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number  
**WO 01/62236 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 31/00**

(21) International Application Number: PCT/US01/03698

(22) International Filing Date: 23 January 2001 (23.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/184,790 24 February 2000 (24.02.2000) US

(71) Applicant (*for all designated States except US*): **PHARMACIA & UPJOHN COMPANY** [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ROGOSKY, Karen** [US/US]; 25 Wycklow Drive, Robbinsville, NJ 08691 (US). **JORN, Deborah** [US/US]; 6 Firethorn Court, Warren, NJ 07059 (US).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(74) Agent: **SOLOMON, Andrew, M.**; Pharmacia & Upjohn Company, Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US).

WO 01/62236 A2

(54) Title: **NEW DRUG COMBINATIONS**

(57) Abstract: A composition comprising: (a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors or a pharmaceutically effective salt thereof; and (b) a pharmaceutically effective amount of one or more antimuscarinic agents or a pharmaceutically effective salt thereof is provided. The composition is useful in treating disorders or diseases of the central nervous system, and particularly useful in treating incontinence.

## NEW DRUG COMBINATIONS

### Background of the Invention

5

#### 1. Field of the Invention

This invention describes new treatments that should provide for a fast acting rapid onset of relief from several nervous system disorders, and it involves the  
10 administration of a norepinephrine reuptake inhibitor, preferably a selective norepinephrine reuptake inhibitor, most preferably the drug reboxetine, in combination with an antimuscarinic agent, preferably tolterodine. In particular, the combination is to be used to treat incontinence.

#### 15 2. Technology Description

The introduction of tricyclic antidepressants in the early 1960s has provided a major advance in the treatment of neuropsychiatric disorders. Reactive and endogenous  
20 depressions, diagnoses formerly carrying grave prognostic implications, have become, with the introduction of the tricyclics, manageable disorders with a much smaller toll on the patient and the society as a whole.

The early tricyclic compounds were reuptake inhibitors of all the catecholamines released in the synaptic cleft, thus resulting in prolongation and enhancement of the  
25 dopamine (DA), noradrenaline (NA) and serotonin (5-hydroxytryptamine = 5-HT) action. Lack of selectivity also causes undesired side effects particularly on the acetylcholine (especially the muscarinic component), and histamine mediated neurotransmission.

30 Because of these unwanted pharmacodynamic activities, cognitive impairment, sedation, urinary and gastrointestinal tract disturbances, and increased intraocular pressure were limiting factors in the clinical use of these compounds and often

required discontinuation of treatment. Of utmost concern were also the cardiac toxic effects and the proconvulsant activity of this group of drugs.

More recently, selective reuptake inhibitors for serotonin (SSRI) have been introduced  
5 with definite advantages in regard to fewer side effects without loss of efficacy. Fluoxetine is an example of such an inhibitor that has had a great amount of commercial success.

Another class of compounds that has been proposed for use in the treatment of  
10 depression is selective norepinephrine reuptake inhibitors. Lower-than-normal levels of norepinephrine are associated with a variety of symptoms including lack of energy, motivation, and interest in life. Thus, a normal level of norepinephrine is essential to maintaining drive and capacity for reward. These neurotransmitters travel from the terminal of a neuron across a small gap (*i.e.*, the synaptic cleft) and bind to receptor  
15 molecules on the surface of a second neuron. This binding elicits intracellular changes that initiate or activate a response or change in the postsynaptic neuron. Inactivation occurs primarily by transport (*i.e.*, reuptake) of the neurotransmitter back into the presynaptic neuron. Abnormality in noradrenergic transmission results in various types of depression, mental, behavioral, and neurological disorders attributed  
20 to a variety of symptoms including a lack of energy, motivation, and interest in life. *See generally*, R.J. Baldessarini, "Drugs and the Treatment of Psychiatric Disorders: Depression and Mania" in *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, McGraw-Hill, NY, NY, pp. 432-439 (1996).

25 Examples of norepinephrine reuptake inhibitors (both selective and not selective) include, but are not limited to the following: tandamine (CAS 42408-80-0; US 3904617; US 4118394), pirandamine (CAS 42408-79-7; US 3995052), ciclazindol (CAS 37751-39-6; US 3891644; US 3957819; US 3976645), fluparoxan (US 4880801), lortalamine (CAS 70384-91-7; US 4201783), talsupram (CAS 21489-20-  
30 3), talopram (CAS 7182-51-6), prindamine, nomifensine (US 3577424), viloxazine (US 3712890), tomoxetine (US 4314081), duloxetine (US 5023269), venlafaxine (US 4535186), milnacipran (US 4478836) and reboxetine (US 4229449).

Tomoxetine, (R)-(-)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine is disclosed in U.S. Patent No. 4,314,081. Reboxetine, 2-[alpha-(2-ethoxy)phenoxybenzyl]morpholine is disclosed in U.S. Patent No. 4,229,449. Reboxetine includes both the racemate, as well as the (-)(R,R) and (+)(S,S) enantiomers. This product is also identified by the following trademarks: VESTRA, PROLIFT, NOREBOX and ERDONAX. Duloxetine, N-methyl-3-(1-naphthalenyloxy)-3-(2-thienyl) propanamine is disclosed in U.S. Patent No. 4,956,388. It is usually administered as the hydrochloride salt and as the (+) enantiomer. Venlafaxine is identified as Compound A of U.S. Patent No. 4,761,501. Milnacipran, N,N-diethyl-2-aminomethyl-1-phenylcyclopropanecarboxamide is disclosed in U.S. Patent No. 4,478,836. To the extent necessary for completion, these above-cited documents are expressly incorporated by reference.

Antimuscarinic agents can be used to treat urinary incontinence. Examples of antimuscarinic agents include, but are not limited to the following: tolterodine, propiverine, oxybutynin, trospium, darifenacin, temiverine, ipratropium.

Tolterodine, Phenol, 2-[3-[bis(1-methylethyl)amino]-1-phenylpropyl]-4-methyl-, (R)-, an antimuscarinic with a high degree of bladder selectivity, has been developed and launched by Pharmacia & Upjohn under the DETROL® trademark for the treatment of incontinence associated with an overactive bladder. This product is disclosed in U.S. Patent No. 5,382,600. The active metabolites of tolterodine are

Propiverine is 1-methyl-4-piperidyl .alpha.,.alpha.-diphenyl-.alpha.-(n-propoxy)acetate and is disclosed in East German Patent No. 106643 and in CAS 82-155841s (1975). Oxybutynin is 4-(Diethylamino)-2-butynylalpha-phenylcyclohexaneglycolate and is disclosed in UK Patent No. 940540. Trospium is 3alpha-Hydroxyspiro[1alphaH,5alphaH-nortropane-8,1'-pyrrolidinium]chloride benzilate and is disclosed in U.S. Patent No. 3480623. Darifenacin is 3-Pyrrolidineacetamide, 1-[2-(2,3-dihydro-5-benzofuranyl)ethyl]-alpha, alpha-diphenyl-, and is disclosed in U.S. Patent No. 5,096,890. Temiverine is Benzeneacetic acid, .alpha.-cyclohexyl-.alpha.-hydroxy-, 4-(diethylamino)-1,1-dimethyl-2-butynyl ester

and is disclosed in U.S. Patent No. 5036098 and ipratropium is 8-isopropylnoratropine methobromide and disclosed in U.S. Patent No. 3505337.

Despite the above advances in the art, it would be desirable to develop a  
5 pharmaceutical composition that would have both the benefits of the norepinephrine reuptake inhibitors and that of the antimuscarinic agents.

#### Brief Summary of the Invention

10 In accordance with the present invention a novel pharmaceutical composition is provided. More specifically, the composition combines one or more selective norepinephrine reuptake inhibitors with one or more antimuscarinic agents, preferably tolterodine. The composition is considered to be particularly effective against incontinence in general, and more particularly stress incontinence.

15

A first embodiment of the present invention provides a composition comprising:

(a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors or a pharmaceutically effective salt thereof; and

20

(b) a pharmaceutically effective amount of one or more antimuscarinic agents or a pharmaceutically effective salt thereof.

In particularly preferred embodiments, component (a) comprises reboxetine in either its  
25 enantiomeric or racemic form and component (b) comprises tolterodine, including its active metabolites.

Yet another embodiment of the present invention provides a method for treating or preventing incontinence or diseases or disorders of the central nervous system  
30 comprising administering a therapeutically effective amount of the above composition to a mammal. In most instances, the mammal will be a human, and the disease or disorder to be treated is incontinence.

A further embodiment of the present invention comprises the use of the above composition to prepare a medicament for treating or preventing incontinence or diseases or disorders of the central nervous system.

- 5 An object of the present invention is to provide novel compositions having biological activity.

A further object of the present invention is to provide a method for treating or preventing incontinence or diseases of the central nervous system by using the novel  
10 compositions of the present invention.

An additional object of the present invention is to provide an effective treatment for incontinence.

- 15 These, and other objects, will readily be apparent to those skilled in the art as reference is made to the detailed description of the preferred embodiment.

#### Detailed Description of the Preferred Embodiment

- 20 In describing the preferred embodiment, certain terminology will be utilized for the sake of clarity. Such terminology is intended to encompass the recited embodiment, as well as all technical equivalents which operate in a similar manner for a similar purpose to achieve a similar result. To the extent that any pharmaceutically active compound is disclosed or claimed, it is expressly intended to include all active metabolites produced  
25 *in vivo*, and, is expressly intended to include all enantiomers, isomers or tautomers where the compound is capable of being present in its enantiomeric, isomeric or tautomeric form.

- The present invention provides a novel composition which is a combination of different  
30 chemical entities, more specifically, the first entity being a norepinephrine reuptake inhibitor and the second being an antimuscarinic agent.

The first component is a norepinephrine reuptake inhibitor, with selective norepinephrine reuptake inhibitors being particularly preferred. This list of compounds includes, but is not limited to the following: tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, 5 viloxazine, tomoxetine, duloxetine, venlafaxine, milnacipran and reboxetine, with reboxetine being particularly preferred.

Examples of pharmaceutically effective salts for the selective norepinephrine reuptake inhibitor include, but are not limited to salts prepared from pharmaceutically 10 acceptable acids or bases, including organic and inorganic acids and bases. When the preferred compound of use is basic (for example reboxetine), salts may be prepared from pharmaceutically acceptable acids. Suitable pharmaceutically acceptable acids include acetic, benzenesulfonic (besylate), benzoic, p-bromophenylsulfonic, camphorsulfonic, carbonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, 15 hydrobromic, hydrochloric, hydroiodic, isethionic, lactic, maleic, malic, mandelic, methanesulfonic (mesylate), mucic, nitric, oxalic, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic, and the like. Examples of such pharmaceutically acceptable salts include, but are not limited to, acetate, benzoate, hydroxybutyrate, bisulfate, bisulfite, bromide, butyne-1,4-dioate, carpoate, chloride, 20 chlorobenzoate, citrate, dihydrogenphosphate, dinitrobenzoate, fumarate, glycollate, heptanoate, hexyne-1,6-dioate, hydroxybenzoate, iodide, lactate, maleate, malonate, mandelate, metaphosphate, methanesulfonate, methoxybenzoate, methylbenzoate, monohydrogenphosphate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, oxalate, phenylbutyrate, phenylproionate, phosphate, phthalate, phylacetate, propanesulfonate, 25 propiolate, propionate, pyrophosphate, pyrosulfate, sebacate, suberate, succinate, sulfate, sulfite, sulfonate, tartrate, xylenesulfonate, and the like.

In particularly preferred embodiments the selective norepinephrine reuptake inhibitor is reboxetine, 2-[ $\alpha$ -((2-ethoxyphenoxy)benzyl)-morpholine, and its pharmaceutically 30 acceptable salts, in either its enantiomeric (particularly the (S,S) enantiomer) or racemic form. Synthesis of racemic reboxetine is described in greater detail in U.S. Patent No. 4,229,449. Individual stereoisomers of reboxetine can be obtained by resolution of the racemic mixture of enantiomers using conventional methods

generally known by those skilled in the art. Such methods include, but are not limited to, resolution by simple crystallization and chromatographic techniques, for example, as set forth in GB 2,167,407. Other methods of preparation are described in US 5,068,433 and US 5,391,735. Reboxetine can be a free base form, or it can be in salt form, preferably the methanesulfonate salt (also called reboxetine mesylate). To the extent necessary for completion, the above patents are expressly incorporated by reference.

The selection of the dosage of the first component is that which can provide relief to the patient. As is well known, the dosage of this component depends on several factors such as the potency of the selected specific compound, the mode of administration, the age and weight of the patient, the severity of the condition to be treated, and the like. This is considered to be within the skill of the artisan and one can review the existing literature on the components to determine optimal dosing. To the extent necessary for completion, the synthesis of the components and dosages described in the patents or CAS documents referenced in the Technology Description portion of this document are expressly incorporated by reference

Desirably, when reboxetine is selected as the active agent, the daily dose contains from about 0.1mg. to about 10 mg. More preferably, each dose of the component contains about 0.5 to about 8 mg of the active ingredient, and even more preferably, each dose contains from about 0.5 to about 5 mg of the active ingredient. This dosage form permits the full daily dosage to be administered in one or two oral doses. This will allow for final formulations containing 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, or 2.5 mg of active. More than once daily or twice daily administrations (e.g., 3, 4, 5 or 6 administrations per day) are also expressly contemplated herein.

The average daily adult dosage of the other norepinephrine reuptake inhibitors is as follows. The dosages expressly include all numerical values, whole or fractional, within the stated range. Pediatric dosages may be less.

Component	Average Daily Dosage (mg/day/patient)
-----------	---------------------------------------



Tandamine	7.5 to 3750
Pirandamine	7.5 to 3750
Ciclazindol	5 to 500
Fluparoxan	.75 to 750
Lortalamine	1 to 200
Talsupram	1 to 3750
Talopram	1 to 3750
Prindamine	1 to 3750
Nomifensine	1 to 80
Viloxazine	1 to 3750
Tomoxetine	1 to 200
Duloxetine	5 to 500
Venlafaxine	2 to 200
Milnacipran	7.5 to 75

The second component comprises one or more antimuscarinic agents. Examples of such agents include tolterodine, propiverine, oxybutynin, trospium, darifenacin, temiverine and ipratropium. Particularly preferred is tolterodine.

5

The chemical name of tolterodine is Phenol, 2-[3-[bis(1-methylethyl)amino]-1-phenylpropyl]-4-methyl-, (R)-, including its pharmaceutically active salts, such as those described above with respect to the first active component and its active metabolites that are produced *in vivo*. Synthesis of tolterodine is disclosed in U.S.

10 Patent No. 5,382,600. To the extent necessary for completion, this patent is expressly incorporated by reference. This compound is particularly useful as an anticholinergic agent, and more specifically for the treatment of incontinence.

15 As is well known, the dosage and administrative regimen (i.e., one, two, three or more administrations per day) of the second component depends on the factors referred to in connection with the dosage selection of the first component. To the extent necessary for completion, the synthesis of the components and dosages described in the patents or CAS documents referenced in the Technology Description portion of this document are expressly incorporated by reference. The average adult daily dosage of the second

component is from about 0.05 mg to about 5 mg per kilogram of body weight, administered in one or more doses, e.g. containing from about 0.05 to about 250 mg each. The dosages expressly include all numerical values, whole or fractional, within the stated range. Pediatric dosages may be less.

5

Compositions of the present invention can conveniently be administered in a pharmaceutical composition containing the active components in combination with a suitable excipient. Such pharmaceutical compositions can be prepared by methods and contain excipients which are well known in the art. A generally recognized compendium of such methods and ingredients is Remington's Pharmaceutical Sciences by E.W. Martin (Mark Publ. Co., 15th Ed., 1975). To the extent necessary for completion, this reference is hereby incorporated by reference. The compositions of the present invention can be administered parenterally (for example, by intravenous, intraperitoneal, subcutaneous or intramuscular injection), topically, orally, intranasally, intravaginally, or rectally, with oral administration being particularly preferred.

For oral therapeutic administration, the inventive composition may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums, foods and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 0.1 to about 100% of the weight of a given unit dosage form. The amount of active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. The above listing is merely representative and one skilled in the art could envision other binders, excipients, sweetening agents and

the like. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules  
5 may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In  
10 addition, the active components may be incorporated into sustained-release preparations and devices including, but not limited to, those relying on osmotic pressures to obtain a desired release profile. Once daily formulations for each of the active components are specifically included.

15 The inventive composition, containing the two active components, may be administered in the same physical form or concomitantly according to the above-described dosages and in the above-described delivery vehicles. The dosages for each active component can be measured separately and can be given as a single combined dose or given separately. They may be given at the same or at different times as long  
20 as both actives are in the patient at one time over a 24-hour period. Concomitant or concurrent administration means the patient takes one drug within about 5 minutes of taking the other drug. Because the goal is to provide rapid symptomatic relief to the patient, in most cases when treatment is started the two drugs would be administered to the patient close in time and typically concomitantly; thereafter, the timing of each  
25 drug's administration may not be as important.

The inventive composition is used to treat incontinence or any of the diseases or disorders of the central nervous system. Such diseases and disorders are defined in The Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV) (American  
30 Psychiatric Association (1995)). To the extent necessary for completion, the contents of this reference and all of the defined diseases or disorders are expressly incorporated by reference. Also considered is the treatment of incontinence of any type. Representative diseases or disorders include, but are not limited to the following:

obesity, depression, schizophrenia, a stress related disease (e.g. general anxiety disorder), panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, incontinence, a stress induced problem with the urinary, gastrointestinal or cardiovascular system (e.g., stress incontinence),  
5 neurodegenerative disorders, autism, chemotherapy-induced vomiting, hypertension, migraine headaches, cluster headaches, sexual dysfunction in a mammal (e.g. a human), addictive disorder and withdrawal syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to general medical conditions, attention-deficit hyperactivity disorder,  
10 bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, a movement disorder (e.g., Tourette's syndrome), oppositional defiant disorder, a pain disorder, peripheral neuropathy, post-traumatic stress disorder,  
15 premenstrual dysphoric disorder, a psychotic disorder, seasonal affective disorder, a sleep disorder, a specific developmental disorder, and selective serotonin reuptake inhibition (SSRI) "poop out" syndrome. Treatment of the above diseases or disorders is accomplished by delivering a therapeutically effective amount of the inventive composition to a mammal. In most cases this will be a human being, but treatment of  
20 food animals (e.g., livestock and poultry) and companion animals (e.g., dogs, cats and horses) is expressly covered herein.

In particular, the inventive composition is to be used in the treatment of incontinence (i.e., stress incontinence, genuine stress incontinence, and mixed incontinence). Stress  
25 urinary incontinence is a symptom describing involuntary loss of urine on carrying out any activity that raises intra-abdominal pressure such as coughing or sneezing. Stress incontinence is also a clinical sign, that is the observation by a care giver of a jet of urine escaping from the urethral meatus (opening) when the patient coughs or strains. Genuine Stress Incontinence (urge incontinence) is the pathological diagnosis of an  
30 incompetent urethral sphincter as diagnosed by Urodynamic testing. Mixed incontinence is stress incontinence in combination with urge incontinence. The latter is a part of the symptom complex of the Overactive Bladder. Retention may be due to outflow obstruction (e.g., high urethral pressure), poor detrusor (bladder muscle)

contractility or lack of coordination between detrusor contraction and urethral relaxation. The inventive drug combination can be used in connection with stress incontinence, urge incontinence or mixed incontinence.

- 5 The novel composition is expected to provide rapid relief to those suffering from the above diseases or disorders with a minimal amount of deleterious side effects.

The invention is described in greater detail by the following non-limiting example.

10

#### Example 1

A pharmaceutical composition is prepared by combining reboxetine in either its racemic or (S,S) enantiomeric form with tolterodine in a pharmaceutically acceptable carrier. The composition contains respective amounts of reboxetine and tolterodine  
15 to deliver on a daily basis between about 0.1 mg to about 10 mg reboxetine and between about 0.05 mg to about 4 mg of tolterodine per kilogram of patient body weight (for example, 3 mg to 240 mg tolterodine for a person weighing 60 kg). The composition is administered to a patient for the treatment of incontinence, and particularly stress incontinence, urge incontinence or mixed incontinence.

20

#### Example 2

A first pharmaceutical composition is prepared by combining reboxetine in either its racemic or +(S,S) enantiomeric form in a pharmaceutically acceptable carrier such  
25 that it can deliver between about 0.1 mg to about 10 mg reboxetine on a daily basis. A second pharmaceutical composition is prepared by combining tolterodine in a pharmaceutically acceptable carrier such that it can deliver between about 0.05 mg to about 4 mg of tolterodine per kilogram of patient body weight on a daily basis. The first composition is administered to a patient suffering from one or more forms of  
30 incontinence once, twice, three times, four times or six times daily such that the daily dosage is between about 0.1 to about 10 mg. The second composition is administered to the same patient at the same time as the administration of the first composition or any time within 24 hours of the administration of the first composition once, twice,

three times, four times or six times daily such that the daily dosage is between about 0.05 mg to about 4 mg of tolterodine per kilogram of patient body weight. Alternatively, the second composition could first be administered, followed by the administration of the first composition as disclosed at the same time, or within 24  
5 hours thereof.

Having described the invention in detail and by reference to the preferred embodiments thereof, it will be apparent that modifications and variations are possible without departing from the scope of the appended claims.

10

What is claimed is:

1. A composition comprising:

5

(a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors or a pharmaceutically effective salt thereof; and

10 (b) a pharmaceutically effective amount of one or more antimuscarinic agents or a pharmaceutically effective salt thereof.

2. The composition according to claim 1 wherein component (a) is selected from the group consisting of tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine,  
15 venlafaxine, milnacipran and reboxetine and mixtures thereof and wherein component (b) is selected from the group consisting of tolterodine, propiverine, oxybutynin, trospium, darifenacin, temiverine and ipratropium and mixtures thereof.

3. The composition according to claim 2 wherein component (a) is reboxetine in  
20 either its racemic or +(S,S) enantiomeric form and component (b) is tolterodine, its active metabolites or mixtures thereof.

4. The composition according to claim 3 containing between about 0.1 mg to about 10 mg reboxetine and between about 0.05 mg to about 4 mg of tolterodine per  
25 kilogram of patient body weight.

5. The composition according to claim 1 wherein component (a) and component (b) are maintained in the same delivery vehicle.

30 6. The composition according to claim 1 wherein component (a) and component (b) are maintained in different delivery vehicles.

7. A method for treating incontinence or a disease or disorder of the central nervous system in a mammal comprising administering to said mammal a pharmaceutically effective amount of a composition comprising:
- 5 (a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors or a pharmaceutically effective salt thereof; and
- (b) a pharmaceutically effective amount of one or more antimuscarinic agents or a pharmaceutically effective salt thereof.
- 10
8. The method according to claim 7 wherein said disease or disorder is selected from the group consisting of obesity, depression, schizophrenia, a stress related disease (e.g. general anxiety disorder), panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, incontinence, a stress
- 15 induced problem with the urinary, gastrointestinal or cardiovascular system, neurodegenerative disorders, autism, chemotherapy-induced vomiting, hypertension, migraine headaches, cluster headaches, sexual dysfunction in a mammal, addictive disorder and withdrawal syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to
- 20 general medical conditions, attention-deficit hyperactivity disorder, bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, a movement disorder, oppositional defiant disorder, a pain disorder, peripheral neuropathy, post-traumatic
- 25 stress disorder, premenstrual dysphoric disorder, a psychotic disorder, seasonal affective disorder, a sleep disorder, a specific developmental disorder, and selective serotonin reuptake inhibition (SSRI) "poop out" syndrome.
9. The method of claim 7 wherein said composition is administered rectally,
- 30 topically, orally, sublingually, intranasally, transdermally or parenterally.
10. The method of claim 7 wherein component (a) and component (b) of said composition are simultaneously administered.



11. The method of claim 7 wherein component (a) and component (b) of said composition are concomitantly administered.
- 5 12. The method according to claim 7 wherein said disease or disorder comprises incontinence.
13. The method according to claim 12 wherein said incontinence is stress incontinence, genuine stress incontinence or mixed incontinence.
- 10 14. The method according to claim 13 wherein component (a) of said composition comprises reboxetine in its racemic or enantiomeric form and component (b) of said composition comprises tolterodine, its active metabolites or mixtures thereof.
- 15 15. The method according to claim 12 wherein said composition contains between about 0.1 mg to about 10 mg reboxetine and between about 0.05 mg to about 4 mg of tolterodine per kilogram of patient body weight.
16. A composition consisting essentially of:
- 20 (a) a pharmaceutically effective amount of reboxetine in its racemic or enantiomeric form or a pharmaceutically effective salt thereof; and
- (b) a pharmaceutically effective amount of tolterodine, its active metabolites or combinations thereof or a pharmaceutically effective salt thereof;
- 25 wherein components (a) and (b) are maintained in the same or in different delivery vehicles.
- 30 17. The use of a composition comprising:
- (a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors or a pharmaceutically effective salt thereof; and

(b) a pharmaceutically effective amount of one or more antimuscarinic agents or a pharmaceutically effective salt thereof to prepare a medicament for treating or preventing incontinence or diseases or disorders of the central nervous system.

5

18. The use according to claim 17 wherein component (a) comprises reboxetine in its racemic or enantiomeric form and component (b) comprises tolterodine, its active metabolites or mixtures thereof.

10 19. A composition comprising:

(a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors or a pharmaceutically effective salt thereof; and

15 (b) a pharmaceutically effective amount of one or more antimuscarinic agents or a pharmaceutically effective salt thereof for use as a medicament.

20. A composition comprising:

20 (a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors selected from the group consisting of tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxctine, duloxetine, venlafaxine, milnacipran and reboxetine and mixtures thereof or a pharmaceutically effective salt thereof; and

25

(b) a pharmaceutically effective amount of one or more antimuscarinic agents selected from the group consisting of tolterodine, propiverine, oxybutynin, trospium, darifenacin, temiverine and ipratropium and mixtures thereof or a pharmaceutically effective salt thereof;

30

21. The composition according to claim 20 wherein component (a) is reboxetine in either its racemic or +(S,S) enantiomeric form and component (b) is darifenacin.

22. The composition according to claim 20 wherein component (a) is reboxetine in either its racemic or +(S,S) enantiomeric form and component (b) is trospium.
23. The composition according to claim 20 wherein component (a) is reboxetine in  
5 either its racemic or +(S,S) enantiomeric form and component (b) is ipratropium.
24. The composition according to claim 20 wherein component (a) is duloxetine.
25. The composition according to claim 23 wherein component (b) is selected from  
10 the group consisting of tolterodine, darifenacin, trospium, ipratropium and mixtures thereof.
26. The composition according to claim 25 containing between about 1 mg to about 200 mg duloxetine and between about 0.05 mg to about 5 mg of component (b) per  
15 kilogram of patient body weight.
27. The composition according to claim 20 wherein component (a) and component (b) are maintained in the same delivery vehicle.
- 20 28. The composition according to claim 20 wherein component (a) and component (b) are maintained in different delivery vehicles.
29. A method for treating incontinence or a disease or disorder of the central nervous system in a mammal comprising administering to said mammal a  
25 pharmaceutically effective amount of a composition comprising:
- (a) a pharmaceutically effective amount of one or more norepinephrine reuptake inhibitors tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine,  
30 milnacipran and reboxetine and mixtures thereof or a pharmaceutically effective salt thereof; and

(b) a pharmaceutically effective amount of one or more antimuscarinic agents selected from the group consisting of tolterodine, propiverine, oxybutynin, trospium, darifenacin, temiverine and ipratropium and mixtures thereof or a pharmaceutically effective salt thereof.

5

30. The method according to claim 29 wherein said disease or disorder is selected from the group consisting of obesity, depression, schizophrenia, a stress related disease (e.g. general anxiety disorder), panic disorder, a phobia, obsessive compulsive disorder, post-traumatic-stress syndrome, immune system depression, incontinence, a stress induced problem with the urinary, gastrointestinal or cardiovascular system, neurodegenerative disorders, autism, chemotherapy-induced vomiting, hypertension, migraine headaches, cluster headaches, sexual dysfunction in a mammal, addictive disorder and withdrawal syndrome, an adjustment disorder, an age-associated learning and mental disorder, anorexia nervosa, apathy, an attention-deficit disorder due to  
15 general medical conditions, attention-deficit hyperactivity disorder, bipolar disorder, bulimia nervosa, chronic fatigue syndrome, conduct disorder, cyclothymic disorder, dysthymic disorder, fibromyalgia and other somatoform disorders, generalized anxiety disorder, an inhalation disorder, an intoxication disorder, a movement disorder, oppositional defiant disorder, a pain disorder, peripheral neuropathy, post-traumatic stress disorder, premenstrual dysphoric disorder, a psychotic disorder, seasonal  
20 affective disorder, a sleep disorder, a specific developmental disorder, and selective serotonin reuptake inhibition (SSRI) "poop out" syndrome.

31. The method of claim 30 wherein component (a) and component (b) of said  
25 composition are simultaneously administered.

32. The method of claim 30 wherein component (a) and component (b) of said composition are concomitantly administered.

30 33. The method according to claim 30 wherein said disease or disorder comprises incontinence.

34. The method according to claim 33 wherein said incontinence is stress incontinence, genuine stress incontinence or mixed incontinence.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number  
**WO 01/62236 A3**

(51) International Patent Classification<sup>7</sup>: A61K 31/535,  
31:135, A61P 13/10, A61K 45/06 // (A61K 31/535,  
31:135)

(21) International Application Number: PCT/US01/03698

(22) International Filing Date: 23 January 2001 (23.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/184,790 24 February 2000 (24.02.2000) US

(71) Applicant (for all designated States except US): PHAR-  
MACIA & UPJOHN COMPANY [US/US]; 301 Henri-  
etta Street, Kalamazoo, MI 49001 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ROGOSKY, Karen  
[US/US]; 25 Wycklow Drive, Robbinsville, NJ 08691  
(US). JORN, Deborah [US/US]; 6 Firethorn Court,  
Warren, NJ 07059 (US).

(74) Agent: SOLOMON, Andrew, M.; Pharmacia & Upjohn  
Company, Intellectual Property Legal Services, 301 Henri-  
etta Street, Kalamazoo, MI 49001 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
7 March 2002

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NEW DRUG COMBINATIONS COMPRISING A NOREPINEPHRINE REUPTAKE INHIBITOR AND AN  
ANTIMUSCARINIC AGENT

(57) Abstract: A composition comprising: (a) a pharmaceutically effective amount of one or more norepinephrine reuptake in-  
hibitors or a pharmaceutically effective salt thereof; and (b) a pharmaceutically effective amount of one or more antimuscarinic  
agents or a pharmaceutically effective salt thereof is provided. The composition is useful in treating disorders or diseases of the  
central nervous system, and particularly useful in treating incontinence.

WO 01/62236 A3



# INTERNATIONAL SEARCH REPORT

Intern. of Application No

PCT/US 01/03698

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/535 A61K31/135 A61P13/10 A61K45/06  
 //(A61K31/535, 31:135)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, BIOSIS, EMBASE, CHEM ABS Data, PHARMAPROJECTS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 368 682 A (EURO CELTIQUE SA) 16 May 1990 (1990-05-16) *cf. abstract, page 2, lines 1-23* ---	1-34
A	WO 97 09036 A (KNOTT TREVOR JOHN ; LESLIE STEWART THOMAS (GB); BROWN ADRIAN (GB);) 13 March 1997 (1997-03-13) *cf. abstract, page 2, para. 7 bridging with page 3, items "vi." and "x."* ---	1-34
A	PHARMACIA & UPJOHN: "pharmacia & upjohn" FORMULARY'S RESOURCE GUIDE, vol. 34, no. 10 suppl., October 1999 (1999-10), pages 74-78, XP001013175 * cf. page 75, right col., last para. bridging with page 76, left col., page 77, table 2* -----	1-34

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

9 August 2001

Date of mailing of the international search report

13/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Stoltner, A

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-6, 17-19 in part

Present claims 1-6 and 19, relate to a product defined by functional terms

by reference to the following parameter(s):

P1: norepinephrine reuptake inhibitors

P2: antimuscarinic agents

The use of these parameters in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. It is impossible to compare the parameters the applicant has chosen to employ with what is set out in the prior art. The lack of clarity is such as to render a meaningful complete search impossible. Consequently, the search has been restricted to the compounds according to claims 2,3,16 and 20.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern .al Application No

PCT/US 01/03698

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0368682 A	16-05-1990	AT 123218 T	15-06-1995
		AU 621949 B	26-03-1992
		AU 4453089 A	17-05-1990
		CA 2002492 A	11-05-1990
		DE 68922894 D	06-07-1995
		DE 68922894 T	11-01-1996
		DK 564589 A	12-05-1990
		ES 2075061 T	01-10-1995
		GB 2225941 A, B	20-06-1990
		IE 68241 L	11-05-1990
		JP 2243618 A	27-09-1990
		JP 2825202 B	18-11-1998
		US 5071646 A	10-12-1991
		ZA 8908431 A	25-07-1990
WO 9709036 A	13-03-1997	AU 6834796 A	27-03-1997
		EP 0847273 A	17-06-1998
		US 6077532 A	20-06-2000